

POSTNIKOV PIECES AND $B\mathbb{Z}/p$ -HOMOTOPY THEORY

NATÀLIA CASTELLANA, JUAN A. CRESPO, AND JÉRÔME SCHERER

ABSTRACT. We present a constructive method to compute the cellularization with respect to $B^m\mathbb{Z}/p$ for any integer $m \geq 1$ of a large class of H -spaces, namely all those which have a finite number of non-trivial $B^m\mathbb{Z}/p$ -homotopy groups (the pointed mapping space $\text{map}_*(B^m\mathbb{Z}/p, X)$ is a Postnikov piece). We prove in particular that the $B^m\mathbb{Z}/p$ -cellularization of an H -space having a finite number of $B^m\mathbb{Z}/p$ -homotopy groups is a p -torsion Postnikov piece. Along the way, we characterize the $B\mathbb{Z}/p^r$ -cellular classifying spaces of nilpotent groups.

INTRODUCTION

The notion of A -homotopy theory was introduced by Dror Farjoun [9] for an arbitrary connected space A . Here A and its suspensions play the role of the spheres in classical homotopy theory, and so the A -homotopy groups of a space X are defined to be the homotopy classes of pointed maps $[\Sigma^i A, X]$. The analogue to weakly contractible spaces are those spaces for which all A -homotopy groups are trivial. This means that the pointed mapping space $\text{map}_*(A, X)$ is contractible, i.e. X is an A -null space. On the other hand, the classical notion of CW -complex is replaced by the one of A -cellular space. Such spaces can be constructed from A by means of pointed homotopy colimits.

Thanks to work of Bousfield [2] and Dror Farjoun [9] there is a functorial way to study X through the eyes of A . The nullification $P_A X$ is the biggest quotient of X which is A -null, and $CW_A X$ is the best A -cellular approximation of the space X . Roughly speaking, $CW_A X$ contains all the transcendent information of the mapping space $\text{map}_*(A, X)$, since the latter is equivalent to $\text{map}_*(A, CW_A X)$. Hence, explicit computation of the cellularization would give access to information about $\text{map}_*(A, X)$. The importance of mapping spaces (in the case $A = B\mathbb{Z}/p$) is well established thank to Miller's solution to the Sullivan conjecture [17] and later work.

While many computations of $P_A X$ are present in the literature, very few computations of $CW_A X$ are available. For instance, Chachólski describes a strategy to compute the cellularization $CW_A X$ in [7]. His method has been successfully

Received by the editors November 26, 2004.

2000 *Mathematics Subject Classification*. Primary 55R35; Secondary 55P60, 55P20, 20F18.

Key words and phrases. Cellularization, H -spaces, Postnikov pieces, nilpotent groups.

All three authors were partially supported by MEC grant MTM2004-06686.

The third author was supported by the program Ramón y Cajal, MEC, Spain.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

applied in some cases (cellularization with respect to Moore spaces [21], $B\mathbb{Z}/p$ -cellularization of classifying spaces of finite groups [10]), but it is in general difficult to apply.

An alternative way to compute $CW_A X$ is the following. The nullification map $l : X \rightarrow P_A X$ provides an equivalence $CW_A X \simeq CW_A \overline{P}_A X$, where, as usual, $\overline{P}_A X$ denotes the homotopy fiber of l . This equivalence gives a strategy when $\overline{P}_A X$ is known. Assume for example that X is A -null. Then $\overline{P}_A X$ is contractible, and thus, so is $CW_A X$. From the A -homotopy point of view, the next case in which the A -cellularization should be accessible is when X has only a finite number of A -homotopy groups, that is, some iterated loop space $\Omega^n X$ is A -null. Natural examples of spaces satisfying this condition are the n -connected covers of A -null spaces.

Let us specialize to H -spaces and $A = B^m \mathbb{Z}/p$. Bousfield has determined in [2] the fiber of the nullification map $X \rightarrow P_{B^m \mathbb{Z}/p} X$ when $\Omega^n X$ is $B^m \mathbb{Z}/p$ -null. He shows that, for such an H -space, $\overline{P}_{B^m \mathbb{Z}/p} X$ is a p -torsion Postnikov piece F , whose homotopy groups are concentrated in dimensions from m to $m+n-1$. As F is also an H -space (because l is an H -map), we call it an *H -Postnikov piece*. The cellularization of X (which is again an H -space because CW_A preserves H -structures) therefore coincides with that of a Postnikov piece. In Section 3, we explain how to compute the cellularization of Postnikov pieces, and this enables us to obtain our main result.

Theorem 5.3. *Let X be a connected H -space such that $\Omega^n X$ is $B^m \mathbb{Z}/p$ -null. Then*

$$CW_{B^m \mathbb{Z}/p} X \simeq F \times K(W, m),$$

where F is a p -torsion H -Postnikov piece with homotopy groups concentrated in dimensions from $m+1$ to $m+n-1$ and W is an elementary abelian p -group.

Thus, when X is an H -space with only a finite number of $B^m \mathbb{Z}/p$ -homotopy groups, the cellularization $CW_{B^m \mathbb{Z}/p} X$ is a p -torsion H -Postnikov piece. This is not true in general if we do not assume X to be an H -space. For instance, the $B\mathbb{Z}/p$ -cellularization of $B\Sigma_3$ is a space with infinitely many non-trivial homotopy groups [11]. Also, it is not true for an arbitrary space A that the A -cellularization of an H -space having a finite number of A -homotopy groups is always a Postnikov piece. This fails, for example, when A is the product of the $K(\mathbb{Z}/p, p)$'s, where p runs over the set of all primes, but it could be true for any n -supported p -torsion space A (in the terminology of [2]).

In our previous work [6], we analyzed a large class of H -spaces which fits into the present framework. Namely, if the mod p cohomology of an H -space X is finitely generated as an algebra over the Steenrod algebra, then there must exist an integer n such that $\Omega^n X$ is $B\mathbb{Z}/p$ -null. Hence, we obtain the following.

Proposition 4.2. *Let X be a connected H -space such that $H^*(X; \mathbb{F}_p)$ is finitely generated as an algebra over the Steenrod algebra. Then*

$$CW_{B\mathbb{Z}/p} X \simeq F \times K(W, 1),$$

where F is a 1-connected p -torsion H -Postnikov piece and W is an elementary abelian p -group. Moreover, there exists an integer k such that $CW_{B^m \mathbb{Z}/p} X \simeq *$ for any $m \geq k$.

Our results allow explicit computations which we exemplify by computing in Proposition 4.3 the $B\mathbb{Z}/p$ -cellularization of the n -connected cover of any finite H -space, as well as the $B^m\mathbb{Z}/p$ -cellularizations of the classifying spaces for real and complex vector bundles BU , BO , and their connected covers BSU , BSO , $BSpin$, and $BString$; see Proposition 5.6.

1. A DOUBLE FILTRATION OF THE CATEGORY OF SPACES

As mentioned in the Introduction, the condition that $\Omega^n X$ be $B^m\mathbb{Z}/p$ -null will enable us to compute the $B^m\mathbb{Z}/p$ -cellularization of H -spaces. This section is devoted to giving a picture of how such spaces are related for different choices of m and n .

First of all, we present a lemma which collects various facts that are needed in the rest of the paper.

Lemma 1.1. *Let X be a connected space and $m > 0$. Then,*

- (1) *If X is $B^m\mathbb{Z}/p$ -null, then $\Omega^n X$ is $B^m\mathbb{Z}/p$ -null for all $n \geq 1$.*
- (2) *If X is $B^m\mathbb{Z}/p$ -null, then it is $B^{m+s}\mathbb{Z}/p$ -null for all $s \geq 0$.*
- (3) *If ΩX is $B^m\mathbb{Z}/p$ -null, then X is $B^{m+s}\mathbb{Z}/p$ -null for all $s \geq 1$.*

Proof. For (1), simply apply $\text{map}_*(B\mathbb{Z}/p, -)$ to the path fibration $\Omega X \rightarrow * \rightarrow X$.

Statement (2) is given by Dwyer’s version of Zabrodsky’s lemma [8, Prop. 3.4] applied to the universal fibration $B^m\mathbb{Z}/p \rightarrow * \rightarrow B^{m+1}\mathbb{Z}/p$.

Finally, (3) is proven like (2), using Zabrodsky’s lemma in its connected version [8, Prop. 3.5] (see also Lemma 2.3). Recall that if ΩX is $B^m\mathbb{Z}/p$ -null, then the component $\text{map}(B^m\mathbb{Z}/p, X)_c$ of the constant map is weakly equivalent to X . \square

Of course, the converses of the previous results are not true. For the first statement, take the classifying space of a discrete group at $m = 1$. For the second and third, consider $X = BU$. It is a $B^2\mathbb{Z}/p$ -null space (see Example 1.4), but neither BU nor ΩBU are $B\mathbb{Z}/p$ -null. Observe that in fact $\Omega^n BU$ is never $B\mathbb{Z}/p$ -null. The next result shows that this is the general situation. That is, if a connected space X is $B^{m+1}\mathbb{Z}/p$ -null, then either ΩX is $B^m\mathbb{Z}/p$ -null or none of the iterated loop spaces $\Omega^n X$ is $B^m\mathbb{Z}/p$ -null for $n \geq 1$.

Theorem 1.2. *Let X be a $B^{m+1}\mathbb{Z}/p$ -null space such that $\Omega^k X$ is $B^m\mathbb{Z}/p$ -null for some $k > 0$. Then ΩX is $B^m\mathbb{Z}/p$ -null.*

Proof. It is enough to prove the result for $k = 2$. Consider the fibration


$$K(Q, m + 1) \rightarrow P_{\Sigma^2 B^m\mathbb{Z}/p} X \simeq X \rightarrow P_{\Sigma B^m\mathbb{Z}/p} X,$$

where the fiber is a p -torsion Eilenberg-Mac Lane space by Bousfield’s description of the fiber of the $\Sigma B^m\mathbb{Z}/p$ -nullification [2, Theorem 7.2]. The base space is $B^{m+1}\mathbb{Z}/p$ -null by Lemma 1.1(3) and so is the total space, by assumption. Thus, the pointed mapping space $\text{map}_*(B^{m+1}\mathbb{Z}/p, K(Q, m + 1))$ must be contractible as well, i.e. $Q = 0$. \square

The previous analysis leads to a double filtration of the category of spaces. Let $n \geq 0$ and $m \geq 1$. We introduce the notation

$$\mathcal{S}_m^n = \{X \mid \Omega^n X \text{ is } B^m\mathbb{Z}/p\text{-null}\}.$$

then Lemma 1.1 yields a diagram of inclusions:



Example 1.3. We give examples of spaces in every stage of the filtration.

- (1) \mathcal{S}_1^0 are the spaces that are $B\mathbb{Z}/p$ -null. This contains in particular any finite space (by Miller’s theorem [17, Thm. A]), and, for a nilpotent space X (of finite type with finite fundamental group), to be $B\mathbb{Z}/p$ -null is equivalent to its cohomology $H^*(X; \mathbb{F}_p)$ being locally finite by [22, Corollary 8.6.2].
- (2) If $X\langle n \rangle$ denotes the n -connected cover of a space X , then the homotopy fiber of $\Omega^{n-1}X\langle n \rangle \rightarrow \Omega^{n-1}X$ is a discrete space. Hence, if $X \in \mathcal{S}_m^0$, then $X\langle n \rangle \in \mathcal{S}_m^{n-1}$.
- (3) Observe that $\mathcal{S}_m^n \subset \mathcal{S}_{m+k}^{n-k}$ for all $0 \leq k \leq n$.
- (4) The previous examples provide spaces in every stage of the double filtration. Consider a finite space. It is automatically $B\mathbb{Z}/p$ -null and its n -connected cover $X\langle n \rangle$ lies in \mathcal{S}_1^{n-1} , hence also in $\mathcal{S}_{k+1}^{n-k-1}$ for all $0 \leq k \leq n$.

The next example provides a number of spaces living in \mathcal{S}_m^0 which do not come from the first row of the filtration. Of course their connected covers will be *new* examples of spaces living in \mathcal{S}_m^n .

Example 1.4. Let E_* be a homology theory. If $\tilde{E}^i(K(\mathbb{Z}/p, m)) = 0$ for all i , then the spaces E^i representing the corresponding homology theory are $B^m\mathbb{Z}/p$ -null. If $\tilde{E}^j(K(\mathbb{Z}/p, m-1)) \neq 0$ for some j , then E^j is not $B^{m-1}\mathbb{Z}/p$ -null. In particular, if E_* is periodic, it follows that the spaces E^i are $B^m\mathbb{Z}/p$ -null for all i , but none of their iterated loops are $B^{m-1}\mathbb{Z}/p$ -null.

A first example of such behavior is obtained from complex K-theory: BU is $B^2\mathbb{Z}/p$ -null, but BU and U are not $B\mathbb{Z}/p$ -null (see [18]). Note that real and quaternionic K -theory enjoy the same properties.

For every m , examples of homology theories following this pattern are given by p -torsion homology theories of type III- m as described in [1]. The m th Morava K -theory $K(m)_*$ for p odd is an example of such behavior with respect to Eilenberg-Mac Lane spaces. The spaces representing $K(m)_*$ are $B^{m+1}\mathbb{Z}/p$ -null, but none of their iterated loops are $B^m\mathbb{Z}/p$ -null.

Our aim is to provide tools to compute the $B^m\mathbb{Z}/p$ -cellularization of any H -space lying in the m th row of the above diagram. The key point is the following result of Bousfield [2], which determines the fiber of the nullification map.

Proposition 1.5. *Let $n \geq 0$ and let X be a connected H -space such that $\Omega^n X$ is $B^m\mathbb{Z}/p$ -null. Then there is an H -fibration*

$$F \rightarrow X \rightarrow P_{B^m\mathbb{Z}/p}X,$$

where F is a p -torsion H -Postnikov piece whose homotopy groups are concentrated in dimensions from m to $m + n - 1$. □

Therefore, since $F \rightarrow X$ is a $B^m\mathbb{Z}/p$ -cellular equivalence, we only need to compute the cellularization of a Postnikov piece (which will end up being a Postnikov piece again; see Theorem 3.6). Actually, even more is true.

Proposition 1.6. *Let X be a connected space such that $CW_{B^m\mathbb{Z}/p}X$ is a Postnikov piece. Then there exists an integer n such that $\Omega^n X$ is $B^m\mathbb{Z}/p$ -null.*

Proof. Let us loop once the Chachólski fibration $CW_{B^m\mathbb{Z}/p}X \rightarrow X \rightarrow P_{\Sigma B^m\mathbb{Z}/p}C$ (see [7, Theorem 20.5]). Since $\Omega P_{\Sigma B^m\mathbb{Z}/p}C$ is equivalent to $P_{B^m\mathbb{Z}/p}\Omega C$ by [9, Theorem 3.A.1], we get a fibration over a $B^m\mathbb{Z}/p$ -null base space

$$\Omega CW_{B^m\mathbb{Z}/p}X \rightarrow \Omega X \rightarrow P_{B^m\mathbb{Z}/p}\Omega C.$$

Now there exists an integer n such that $\Omega^n CW_{B^m\mathbb{Z}/p}X$ is discrete, thus $B^m\mathbb{Z}/p$ -null. Therefore, so is $\Omega^n X$. □

2. CELLULARIZATION OF FIBRATIONS OVER BG

In general, it is very difficult to compute the cellularization of the total space of a fibration. In this section, we explain how to deal with this problem when the base space is the classifying space of a discrete group. The first step applies to any group. In the second step (see Proposition 2.4 below) we specialize to nilpotent groups.

Proposition 2.1. *Let $r \geq 1$ and let $F \rightarrow E \xrightarrow{\pi} BG$ be a fibration, where G is a discrete group. Let S be the (normal) subgroup generated by all elements $g \in G$ of order p^i for some $i \leq r$ such that the inclusion $B\langle g \rangle \rightarrow BG$ lifts to E . Then the pullback of the fibration along $BS \rightarrow BG$*

$$\begin{array}{ccccc} E' & \xrightarrow{f} & E & \xrightarrow{p} & B(G/S) \\ \downarrow & & \downarrow \pi & & \parallel \\ BS & \longrightarrow & BG & \xrightarrow{p'} & B(G/S) \end{array}$$

induces a $B\mathbb{Z}/p^r$ -cellular equivalence $f : E' \rightarrow E$ on the total space level.

Proof. We have to show that f induces a homotopy equivalence on pointed mapping spaces $\text{map}_*(B\mathbb{Z}/p^r, -)$. The top fibration in the diagram yields a fibration

$$\text{map}_*(B\mathbb{Z}/p^r, E') \xrightarrow{f_*} \text{map}_*(B\mathbb{Z}/p^r, E) \xrightarrow{p_*} \text{map}_*(B\mathbb{Z}/p^r, B(G/S)).$$

Since the base is homotopically discrete, we only need to check that all components of the total space are sent by p_* to the component of the constant. Thus consider a map $h : B\mathbb{Z}/p^r \rightarrow E$. The composite $p \circ h$ is homotopy equivalent to a map

induced by a group homomorphism $\alpha: \mathbb{Z}/p^r \rightarrow G$ whose image $\alpha(1) = g$ is in S by construction. Therefore $p \circ h = p' \circ \pi \circ h$ is null-homotopic. \square

Remark 2.2. If the fibration in the above proposition is an H -fibration (in particular if G is abelian), the set of elements g for which there is a lift to the total space forms a subgroup of G . The central extension $Z(D_8) \hookrightarrow D_8 \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2$ of the dihedral group D_8 provides an example where the subgroup S is $\mathbb{Z}/2 \times \mathbb{Z}/2$, but the element in S represented by an element of order 4 in D_8 does not admit a lift.

The next lemma is a variation of Dwyer’s version of Zabrodsky’s Lemma in [8].

Lemma 2.3. *Let $F \rightarrow E \xrightarrow{f} B$ be a fibration over a connected base, and let A be a connected space such that ΩA is F -null. Then any map $g: E \rightarrow A$ which is homotopic to the constant map when restricted to the fiber F factors through a map $h: B \rightarrow A$ up to unpointed homotopy and, moreover, g is pointed null-homotopic if and only if h is so.*

Proof. Since ΩA is F -null, we see that the component $\text{map}_*(F, A)_c$ of the constant map is contractible, and therefore, the evaluation at the base point $\text{map}(F, A)_c \rightarrow A$ is an equivalence. By [8, Proposition 3.5], f induces a homotopy equivalence

$$\text{map}(B, A) \simeq \text{map}(E, A)_{[F]},$$

where $\text{map}(E, A)_{[F]}$ denotes the space of maps $E \rightarrow A$ which are homotopic to the constant map when restricted to F .

Looking at the component of the constant map, we see that $\text{map}(B, A)_c \simeq \text{map}(E, A)_c$. Since any map homotopic to the constant map is also homotopic by a pointed homotopy, the result follows. \square

Proposition 2.4. *Let $r \geq 1$ and let $F \xrightarrow{i} E \xrightarrow{\pi} BG$ be a fibration, where G is a nilpotent group generated by elements of order p^i with $i \leq r$. Assume that for each of these generators $x \in G$, the inclusion $B\langle x \rangle \rightarrow BG$ lifts to E . If F is $B\mathbb{Z}/p^r$ -cellular, then so is E .*

Proof. In [7], Chachólski describes the cellularization $CW_{B\mathbb{Z}/p^r} E$ as the homotopy fiber of the composite

$$f: E \rightarrow C \rightarrow P_{\Sigma B\mathbb{Z}/p^r} C,$$

where C is the homotopy cofiber of the evaluation map $\bigvee_{[B\mathbb{Z}/p^r, E]} B\mathbb{Z}/p^r \rightarrow E$. This tells us that E is cellular if the map f is null-homotopic. Observe that if f is null-homotopic, then the fiber inclusion $CW_{B\mathbb{Z}/p^r} E \rightarrow E$ has a section and, therefore, E is cellular, since it is a retract of a cellular space ([9, 2.D.1.5]).

As the existence of an unpointed homotopy to the constant map implies the existence of a pointed one, we now work in the category of unpointed spaces. We remark that for any map $g: Z \rightarrow E$ from a $B\mathbb{Z}/p^r$ -cellular space Z , the composite $f \circ g$ is null-homotopic, since g factors through the cellularization of E . In particular, the composite $f \circ i$ is null-homotopic. By Lemma 2.3, f factors through a map $\bar{f}: BG \rightarrow P_{\Sigma B\mathbb{Z}/p^r} C$ such that $\bar{f} \circ \pi \simeq f$ and, moreover, f is null-homotopic if and only if \bar{f} is so.

We first assume that G is a finite group and show by induction on the order of G that f is null-homotopic. If $|G| = p$, the existence of a section $s: BG \rightarrow E$ implies that $f \circ s = \bar{f}$ is null-homotopic since $BG = B\mathbb{Z}/p$ is cellular.

Let $\{x_1, \dots, x_k\}$ be a minimal set of generators which admit a lift. Let $H \trianglelefteq G$ be the normal subgroup generated by x_1, \dots, x_{k-1} and their conjugates by powers of x_k . There is a short exact sequence

$$H \rightarrow G \rightarrow \mathbb{Z}/p^a,$$

where the quotient group is generated by the image of the generator x_k . Consider the fibration $F \rightarrow E' \rightarrow BH$ obtained by pulling back along $BH \rightarrow BG$, and denote by $h : E' \rightarrow E$ the induced map between the total spaces. Since H satisfies the assumptions of the proposition, the induction hypothesis tells us that E' is cellular and therefore, $f \circ h$ is null-homotopic. This implies that the restriction of \bar{f} to BH is null-homotopic. Consider the following diagram:

$$\begin{array}{ccccc}
 B(\langle x_k \rangle \cap H) & \longrightarrow & BH & & \\
 \downarrow & & \downarrow & \searrow & * \\
 B(\langle x_k \rangle) & \longrightarrow & BG & \xrightarrow{\bar{f}} & P_{\Sigma B\mathbb{Z}/p^r} C \\
 \downarrow & & \downarrow & \nearrow & f' \\
 B\mathbb{Z}/p^a & \xlongequal{\quad} & B\mathbb{Z}/p^a & &
 \end{array}$$

By Lemma 2.3, it is enough to show that f' is null-homotopic. Again, applying Lemma 2.3 to the fibration on the left shows that f' is null-homotopic since \bar{f} restricted to $\langle x_k \rangle$ is so. Therefore, \bar{f} is null-homotopic.

Assume now that G is not finite. Any subgroup of G generated by a finite number of elements of order a power of p has a finite abelianization, and must therefore be itself finite by [20, Theorem 2.26]. Thus, G is locally finite, i.e. G is a filtered colimit of finite nilpotent groups generated by elements of order p^i for $i \leq r$. Likewise, BG is a filtered homotopy colimit of classifying spaces of finite groups (generated by finite subsets of the set of generators) which satisfy the hypotheses of the proposition. The total space E can be obtained as a pointed filtered colimit of the total spaces obtained by pulling back the fibration. By the case when G is finite, these total spaces are all cellular and therefore, so is E . \square

Sometimes the existence of the “local” sections defined for every generator permits the construction of a global section of the fibration. By a result of Chachólski [7, Theorem 4.7], the total space of such a split fibration is cellular since F and BG are so. This is the case for an H -fibration, and E is then weakly equivalent to the product $F \times BG$.

A straightforward consequence of the above proposition (in the case when the fibration is the identity on BG) is the following characterization of the $B\mathbb{Z}/p^r$ -cellular classifying spaces. For $r = 1$, we obtain R. Flores’ result [10, Theorem 4.14].

Corollary 2.5. *Let $r \geq 1$ and let G be a nilpotent group generated by elements of order p^i with $i \leq r$. Then BG is $B\mathbb{Z}/p^r$ -cellular. \square*

Example 2.6. The quaternion group Q_8 of order 8 is generated by elements of order 4. Therefore, BQ_8 is $B\mathbb{Z}/4$ -cellular. We do not know an explicit way to construct BQ_8 as a pointed homotopy colimit of a diagram whose values are copies of $B\mathbb{Z}/4$.

We can now state the main result of this section. It provides a constructive description of the cellularization of the total space of certain fibrations over classifying spaces of nilpotent groups.

Theorem 2.7. *Let G be a nilpotent group and let $F \rightarrow E \rightarrow BG$ be a fibration with $B\mathbb{Z}/p^r$ -cellular fiber F . Then the cellularization of E is the total space of a fibration $F \rightarrow CW_{B\mathbb{Z}/p^r}E \rightarrow BS$, where $S \triangleleft G$ is the (normal) subgroup generated by the p -torsion elements g of order p^i with $i \leq r$, such that the inclusion $B\langle g \rangle \rightarrow BG$ lifts to E .*

Proof. By Proposition 2.1, pulling back along $BS \rightarrow BG$ yields a cellular equivalence f in the following square:

$$\begin{array}{ccc} E_S & \xrightarrow{f} & E \\ \downarrow & & \downarrow \\ BS & \longrightarrow & BG. \end{array}$$

By Proposition 2.4, the total space E_S is cellular and therefore $E_S \simeq CW_{B\mathbb{Z}/p^r}E$. □

Corollary 2.8. *Let G be a nilpotent group and let $S \triangleleft G$ be the (normal) subgroup generated by the p -torsion elements g of order p^i with $i \leq r$. Then $CW_{B\mathbb{Z}/p^r}BG \simeq BS$. Moreover, when G is finitely generated, S is a finite p -group.*

Proof. We only need to show that S is a finite p -group. Note that the abelianization of S is p -torsion. Thus, S is also a torsion group (see [23, Cor. 3.13]). Moreover, since G is finitely generated, S is finite, by [23, 3.10]. □

In fact, Theorem 2.7 also holds when the base space is an Eilenberg-Mac Lane space $K(G, n)$.

Proposition 2.9. *Let n be an integer ≥ 2 and let G be a finitely generated abelian group of exponent dividing p^r . Consider a fibration $F \xrightarrow{i} E \xrightarrow{\pi} K(G, n)$ such that, for each generator $x \in G$, the inclusion $K(\langle x \rangle, n) \rightarrow K(G, n)$ lifts to E . If F is $B\mathbb{Z}/p^r$ -cellular, then so is E . □*

3. CELLULARIZATION OF NILPOTENT POSTNIKOV PIECES

In this section, we compute the cellularization with respect to $B\mathbb{Z}/p^r$ of nilpotent Postnikov pieces. The main difficulty lies in the fundamental group, so it will be no surprise that these results hold as well for cellularization with respect to $B^m\mathbb{Z}/p^r$ with $m \geq 2$. We will often use the following closure property [9, Theorem 2.D.11].

Proposition 3.1. *Let $F \rightarrow E \rightarrow B$ be a fibration where F and E are A -cellular. Then so is B . □*

Example 3.2 ([9, Corollary 3.C.10]). The Eilenberg-Mac Lane space $K(\mathbb{Z}/p^k, n)$ is $B\mathbb{Z}/p^r$ -cellular for any integer k and any $n \geq 2$.

The construction of the cellularization is performed by looking first at the universal cover of the Postnikov piece. We start with the basic building blocks, the Eilenberg-Mac Lane spaces. For the structure results on infinite abelian groups, we refer the reader to Fuchs' book [12].

Lemma 3.3. *An Eilenberg-Mac Lane space $K(A, m)$, with $m \geq 2$, is $B\mathbb{Z}/p^r$ -cellular if and only if A is a p -torsion abelian group.*

Proof. It is clear that A must be p -torsion. Thus, assume that A is a p -torsion group. If A is bounded, it is isomorphic to a direct sum of cyclic groups. Since cellularization commutes with finite products, $K(A, m)$ is $B\mathbb{Z}/p^r$ -cellular when A is a finite direct sum of cyclic groups. By taking a (possibly transfinite) telescope of $B\mathbb{Z}/p^r$ -cellular spaces, we obtain that $K(A, m)$ is $B\mathbb{Z}/p^r$ -cellular for any bounded group.

In general, A splits as a direct sum of a divisible group D and a reduced group T . A p -torsion divisible group is a direct sum of copies of \mathbb{Z}/p^∞ , which is a union of bounded groups. Thus, $K(D, m)$ is cellular. Now T has a basic subgroup $P < T$, which is a direct sum of cyclic groups, and the quotient T/P is divisible. So $K(T, m)$ is the total space of a fibration

$$K(P, m) \rightarrow K(T, m) \rightarrow K(D, m).$$

When $m \geq 3$, we are done because of the closure property Proposition 3.1. If $m = 2$, we have to refine the analysis of the fibration because $K(D, m - 1)$ is not cellular. However, since D is a union of bounded groups $D[p^k]$, the space $K(T, 2)$ is the telescope of total spaces X_k of fibrations with cellular fiber $K(P, 2)$ and base $K(D[p^k], 2)$. We claim that these total spaces are cellular (and thus, so is $K(T, 2)$) and proceed by induction on the bound. Consider the subgroup $D[p^k] < D[p^{k+1}]$ whose quotient is a direct sum of cyclic groups \mathbb{Z}/p . Therefore, X_{k+1} is the base space in a fibration

$$K(\oplus\mathbb{Z}/p, 1) \rightarrow X_k \rightarrow X_{k+1},$$

where the fiber and total space are cellular. We are done. □

We are now ready to prove that any simply connected p -torsion Postnikov piece is a $B\mathbb{Z}/p^r$ -cellular space.

Proposition 3.4. *A simply connected Postnikov piece is $B\mathbb{Z}/p^r$ -cellular if and only if it is p -torsion.*

Proof. Let X be a simply connected p -torsion Postnikov piece. For some integer m , the m -connected cover $X\langle m \rangle$ is an Eilenberg-Mac Lane space, which is cellular by Lemma 3.3. Consider the principal fibration

$$K(\pi_m X, m - 1) \rightarrow X\langle m \rangle \rightarrow X\langle m - 1 \rangle.$$

If $m \geq 3$, both $X\langle m \rangle$ and $K(\pi_m X, m - 1)$ are cellular. It follows that $X\langle m - 1 \rangle$ is cellular by the closure property Proposition 3.1. An iteration of the same argument shows that $X\langle 2 \rangle$ is cellular.

Thus, let us look at the fibration $X\langle 2 \rangle \rightarrow X \rightarrow K(\pi_2 X, 2)$. The discussion in the proof of Lemma 3.3 also applies to the p -torsion group $\pi_2 X$. If this is a bounded group, say of exponent p^k , an induction on the bound shows that X is actually the base space of a fibration where the total space is cellular, because its second homotopy group is of exponent p^{k-1} , and the fiber is cellular because it is of the form $K(V, 1)$, with V a p -torsion abelian group of exponent $\leq p^r$. Then the closure property Proposition 3.1 ensures that X is cellular.

If $\pi_2 X$ is divisible, X is a telescope of cellular spaces, hence cellular. If it is reduced, taking a basic subgroup $B < \pi_2 X$ yields a diagram of fibrations

$$\begin{array}{ccccc}
 X\langle 2 \rangle & \longrightarrow & Y & \longrightarrow & K(B, 2) \\
 \parallel & & \downarrow & & \downarrow \\
 X\langle 2 \rangle & \longrightarrow & X & \longrightarrow & K(\pi_2 X, 2) \\
 \downarrow & & \downarrow & & \downarrow \\
 * & \longrightarrow & K(D, 2) & \xlongequal{\quad} & K(D, 2),
 \end{array}$$

which exhibits X as the total space of a fibration over $K(D, 2)$ with D divisible and a $B\mathbb{Z}/p^r$ -cellular fiber. Therefore, by writing D as a union of bounded groups as in the proof of Lemma 3.3, one obtains X as a telescope of cellular spaces. Thus, X is $B\mathbb{Z}/p^r$ -cellular as well. \square

Remark 3.5. The proof of the proposition holds in the more general setting, where X is a p -torsion space such that $X\langle m \rangle$ is $B\mathbb{Z}/p^r$ -cellular for some $m \geq 2$. The proposition corresponds to the case when some m -connected cover $X\langle m \rangle$ is contractible.

Recall from [13, Corollary 2.12] that a connected space is nilpotent if and only if its Postnikov system admits a principal refinement

$$\cdots \rightarrow X_s \rightarrow X_{s-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0.$$

This means that each map $X_{s+1} \rightarrow X_s$ in the tower is a principal fibration with fiber $K(A_s, i_s - 1)$ for some increasing sequence of integers $i_s \geq 2$. We are only interested in finite Postnikov pieces, i.e. nilpotent spaces that can be constructed in a finite number of steps by taking homotopy fibers of k -invariants $X_s \rightarrow K(A_s, i_s)$.

The key step in the study of the cellularization of a nilpotent finite Postnikov piece is the analysis of principal fibrations (given in our case by the k -invariants).

Theorem 3.6. *Let X be a p -torsion nilpotent Postnikov piece. Then there exists a fibration*

$$X\langle 1 \rangle \rightarrow CW_{B\mathbb{Z}/p^r} X \rightarrow BS,$$

where S is the (normal) subgroup of $\pi_1 X$ generated by the elements g of order p^i with $i \leq r$, such that the inclusion $B\langle g \rangle \rightarrow B\pi_1 X$ admits a lift to X .

Proof. By Proposition 3.4, the universal cover $X\langle 1 \rangle$ is cellular, and there is a fibration $X\langle 1 \rangle \rightarrow X \rightarrow BG$, where $G = \pi_1 X$ is nilpotent. The result then follows from Theorem 2.7. \square

4. CELLULARIZATION OF H -SPACES

In this section, we will use the computations of the cellularization of p -torsion nilpotent Postnikov systems to determine $CW_{B\mathbb{Z}/p} X$ when X is an H -space. We prove:

Theorem 4.1. *Let X be a connected H -space such that $\Omega^n X$ is $B\mathbb{Z}/p$ -null. Then*

$$CW_{B\mathbb{Z}/p} X \simeq Y \times K(W, 1),$$

where Y is a simply connected p -torsion H -Postnikov piece with homotopy groups concentrated in dimensions $\leq n$ and W is an elementary abelian p -group.

Proof. The fibration in Bousfield’s result Proposition 1.5 yields a cellular equivalence between a connected p -torsion H -Postnikov piece F and X . Theorem 3.6 thus applies. Moreover, since F is an H -space as well, the subgroup S is abelian and generated by elements of order p . Therefore, the H -fibration in Theorem 3.6 $F\langle 1 \rangle \rightarrow CW_{B\mathbb{Z}/p}F \rightarrow K(W, 1)$ admits a section (summing up the local sections), and the cellularization splits as a product. \square

This result applies for H -spaces satisfying certain finiteness conditions.

Proposition 4.2. *Let X be a connected H -space such that $H^*(X; \mathbb{F}_p)$ is finitely generated as an algebra over the Steenrod algebra. Then*

$$CW_{B\mathbb{Z}/p}X \simeq F \times K(W, 1),$$

where F is a 1-connected p -torsion H -Postnikov piece and W is an elementary abelian p -group. Moreover, there exists an integer k such that $CW_{B^m\mathbb{Z}/p}X \simeq *$ for any $m \geq k$.

Proof. In [6], we prove that if $H^*(X; \mathbb{F}_p)$ is finitely generated as an algebra over the Steenrod algebra, then $\Omega^n X$ is $B\mathbb{Z}/p$ -null for some $n \geq 0$. Hence, Theorem 4.1 applies, and we obtain the desired result. In addition, Lemma 1.1 shows that X is $B^{n+s+1}\mathbb{Z}/p$ -null for any $s \geq 0$, which implies the second part of the result. \square

The technique we propose in this paper is not only a nice theoretical tool which provides a general statement about what the $B\mathbb{Z}/p$ -cellularization of H -spaces looks like. Our next result shows that one can actually identify this new space precisely when dealing with connected covers of finite H -spaces. Recall that by Miller’s theorem [17, Thm. A], any finite H -space X is $B\mathbb{Z}/p$ -null and hence, $CW_{B\mathbb{Z}/p}X \simeq *$. The universal cover of X is still finite, and thus $CW_{B\mathbb{Z}/p}(X\langle 1 \rangle)$ is contractible as well. We can therefore assume that X is 1-connected. The computation of the cellularization of the 3-connected cover is already implicit in [4].

Proposition 4.3. *Let X be a simply connected finite H -space and let k denote the rank of the free abelian group $\pi_3 X$. Then $CW_{B\mathbb{Z}/p}(X\langle 3 \rangle) \simeq K(\bigoplus_k \mathbb{Z}/p, 1)$. For $n \geq 4$, up to p -completion, the universal cover of $CW_{B\mathbb{Z}/p}(X\langle n \rangle)$ is weakly equivalent to the 2-connected cover of $\Omega(X[n])$.*

Proof. By Browder’s famous result [5, Theorem 6.11], X is even 2-connected and its third homotopy group $\pi_3 X$ is free abelian (of rank k) by Hubbuck and Kane’s theorem [14]. This means we have a fibration

$$K(\bigoplus_k \mathbb{Z}_{p^\infty}, 1) \rightarrow X\langle 3 \rangle \rightarrow P_{B\mathbb{Z}/p}X\langle 3 \rangle,$$

which shows that $CW_{B\mathbb{Z}/p}X\langle 3 \rangle \simeq K(\bigoplus_k \mathbb{Z}/p, 1)$.

We now deal with the higher connected covers. Consider the following commutative diagram of fibrations:

$$\begin{array}{ccccc} F & \xlongequal{\quad\quad\quad} & F & \longrightarrow & * \\ \downarrow & & \downarrow & & \downarrow \\ \Omega X[n] & \longrightarrow & X\langle n \rangle & \longrightarrow & X \\ \downarrow & & \downarrow & & \parallel \\ P_{B\mathbb{Z}/p}(\Omega X[n]) & \longrightarrow & P_{B\mathbb{Z}/p}(x\langle n \rangle) & \longrightarrow & X \end{array}$$

where F is a p -torsion Postnikov piece by [2, Thm 7.2] and the fiber inclusions are all $B\mathbb{Z}/p$ -cellular equivalences, because the base spaces are $B\mathbb{Z}/p$ -null. Therefore,

$$CW_{B\mathbb{Z}/p}(X\langle n \rangle) \simeq CW_{B\mathbb{Z}/p}F \simeq F\langle 1 \rangle \times K(W, 1).$$

We wish to identify $F\langle 1 \rangle$. Since the fibrations in the diagram are nilpotent, by [3, II.4.8] they remain fibrations after p -completion. By Neisendorfer’s theorem [19], the map $P_{B\mathbb{Z}/p}(X\langle n \rangle) \rightarrow X$ is an equivalence up to p -completion, which means that $P_{B\mathbb{Z}/p}(\Omega(X\langle n \rangle))_p^\wedge \simeq *$. Thus $F_p^\wedge \simeq (\Omega(X\langle n \rangle))_p^\wedge$. Note that $\Omega(X\langle n \rangle)$ is simply connected and its second homotopy group is free by the above-mentioned theorem of Hubbuck and Kane (which corresponds up to p -completion to the direct sum of k copies of the Prüfer group \mathbb{Z}/p^∞ in $\pi_1 F$). Hence, $F\langle 1 \rangle$ coincides with $(\Omega(X\langle n \rangle))\langle 2 \rangle$ up to p -completion. \square

To illustrate this result, we compute the $B\mathbb{Z}/2$ -cellularization of the successive connected covers of S^3 . The only delicate point is the identification of the fundamental group.

Example 4.4. Recall that S^3 is $B\mathbb{Z}/2$ -null since it is a finite space. Thus, the cellularization $CW_{B\mathbb{Z}/2}S^3$ is contractible. Next, the fibration

$$K(\mathbb{Z}_{2^\infty}, 1) \rightarrow S^3\langle 3 \rangle \rightarrow P_{B\mathbb{Z}/2}(S^3\langle 3 \rangle)$$

shows that $CW_{B\mathbb{Z}/2}(S^3\langle 3 \rangle) \simeq K(\mathbb{Z}/2, 1)$. Finally, since $S^3\langle 4 \rangle$ does not split as a product (the k -invariant is not trivial), we see that $CW_{B\mathbb{Z}/2}(S^3\langle 4 \rangle) \simeq K(\mathbb{Z}/2, 3)$. Likewise, for any integer $n \geq 4$, we have that $CW_{B\mathbb{Z}/2}(S^3\langle n \rangle)$ is weakly equivalent to the 2-completion of the 2-connected cover of $\Omega(S^3\langle n \rangle)$. The same phenomenon occurs at odd primes.

5. CELLULARIZATION WITH RESPECT TO $B^m\mathbb{Z}/p$

All the techniques developed for fibrations over BG apply to fibrations over $K(G, n)$ when $n > 1$, and we get the following results.

Lemma 5.1. *Let $m \geq 2$ and let X be a connected space. Then $CW_{B^m\mathbb{Z}/p^r}X$ is weakly equivalent to $CW_{B^m\mathbb{Z}/p^r}(X\langle n-1 \rangle)$.*

Proof. Consider the fibrations $X\langle i \rangle \rightarrow X\langle i-1 \rangle \rightarrow K(\pi_i X, i)$. For $i < m$, the base space is $B^m\mathbb{Z}/p^r$ -null and so $CW_{B^m\mathbb{Z}/p^r}(X\langle i \rangle) \simeq CW_{B^m\mathbb{Z}/p^r}(X\langle i-1 \rangle)$. \square

Proposition 5.2. *Let $m \geq 2$ and let X be a p -torsion nilpotent Postnikov piece. Then there exists a fibration*

$$X\langle m \rangle \rightarrow CW_{B^m\mathbb{Z}/p^r}X \rightarrow K(W, m),$$

where W is a p -torsion subgroup of $\pi_m X$ of exponent dividing p^r . \square

Theorem 5.3. *Let X be a connected H -space such that $\Omega^n X$ is $B^m\mathbb{Z}/p$ -null. Then*

$$CW_{B^m\mathbb{Z}/p}X \simeq F \times K(W, m),$$

where F is a p -torsion H -Postnikov piece with homotopy groups concentrated in dimensions from $m+1$ to $m+n-1$, and W is an elementary abelian p -group. \square

Example 5.4. Let X denote “Milgram’s space” (see [16]), the homotopy fiber of $Sq^2 : K(\mathbb{Z}/2, 2) \rightarrow K(\mathbb{Z}/2, 4)$. This is an infinite loop space. By Proposition 3.4, we know it is already $B\mathbb{Z}/2$ -cellular. Since the k -invariant is not trivial, we see that

$$CW_{B^2\mathbb{Z}/2}X \simeq CW_{B^3\mathbb{Z}/2}X \simeq K(\mathbb{Z}/2, 3).$$

Finally, we compute the cellularization of the (infinite loop) space BU and its 2-connected cover BSU with respect to Eilenberg-Mac Lane spaces $B^m\mathbb{Z}/p$. By Bott periodicity, this actually tells us the answer for all connected covers of BU .

Example 5.5. First of all, recall from Example 1.4 that BU is $B^2\mathbb{Z}/p$ -null since $\tilde{K}^*(B^2\mathbb{Z}/p) = 0$ and its iterated loops are never $B\mathbb{Z}/p$ -null. Therefore, the cellularization $CW_{B^m\mathbb{Z}/p}BU$ is contractible if $m \geq 2$. Since $BU \simeq BSU \times BS^1$, the same holds for BSU .

We now compute the $B^m\mathbb{Z}/p$ -cellularization of BO and its connected covers BSO , $BSpin$, and $BString$.

Proposition 5.6. *Let $m \geq 2$. Then*

- (1) $CW_{B^m\mathbb{Z}/p}BO \simeq CW_{B^m\mathbb{Z}/p}BSO \simeq CW_{B^m\mathbb{Z}/p}BSpin \simeq *$,
- (2) $CW_{B^m\mathbb{Z}/p}BString \simeq *$ if $m > 2$,
- (3) $CW_{B^2\mathbb{Z}/p}BString \simeq K(\mathbb{Z}/p, 2)$ and $\text{map}_*(B^2\mathbb{Z}/p, BString) \simeq \mathbb{Z}/p$.

Proof. In [15], W. Meier proves that real and complex K -theory have the same acyclic spaces, hence BO is also $B^2\mathbb{Z}/p$ -null. Therefore, $CW_{B^m\mathbb{Z}/p}BO$ is contractible for any $m \geq 2$. The 2-connected cover of BO is BSO , and there is a splitting $BO \simeq BSO \times B\mathbb{Z}/2$, so that $CW_{B^m\mathbb{Z}/p}BSO \simeq *$.

The 4-connected cover of BO is $BSpin$. From the fibration

$$BSpin \rightarrow BSO \xrightarrow{w_2} K(\mathbb{Z}/2, 2),$$

we infer that the homotopy fiber of $BSpin \rightarrow BSO$ is $B\mathbb{Z}/2$. Since BSO and $B\mathbb{Z}/2$ are $B^2\mathbb{Z}/p$ -null, so is $BSpin$. Therefore, $CW_{B^m\mathbb{Z}/p}BSpin$ is contractible.

Finally, the 8-connected cover of BO is $BString$. It is the homotopy fiber of $BSpin \xrightarrow{p_1/4} K(\mathbb{Z}, 4)$, where p_1 denotes the first Pontrjagin class. Consider the fibration

$$K(\mathbb{Z}, 3) \rightarrow BString \rightarrow BSpin,$$

where the base space is $B^m\mathbb{Z}/p$ -null for $m \geq 2$. Together with the exact sequence $\mathbb{Z} \rightarrow \mathbb{Z}[\frac{1}{p}] \rightarrow \mathbb{Z}/p^\infty$, this implies that

$$CW_{B^m\mathbb{Z}/p}BString \simeq CW_{B^m\mathbb{Z}/p}K(\mathbb{Z}, 3) \simeq CW_{B^m\mathbb{Z}/p}K(\mathbb{Z}/p^\infty, 2).$$

This is a contractible space unless $m = 2$, when we obtain $K(\mathbb{Z}/p, 2)$. The explicit description of the pointed mapping space $\text{map}_*(B^2\mathbb{Z}/p, BString)$ follows. \square

Observe that the iterated loops of the m -connected covers of BO and BU are never $B\mathbb{Z}/p$ -null. Hence, their cellularizations with respect to $B\mathbb{Z}/p$ must have infinitely many non-vanishing homotopy groups by Proposition 1.6.

ACKNOWLEDGEMENTS

We would like to thank Daniel Davis for carefully revising our English.

REFERENCES

- [1] A. K. Bousfield, *Constructions of factorization systems in categories*, J. Pure Appl. Algebra **9** (1976/77), no. 2, 207–220. MR0478159 (57:17648)
- [2] ———, *Localization and periodicity in unstable homotopy theory*, J. Amer. Math. Soc. **7** (1994), no. 4, 831–873. MR1257059 (95c:55010)
- [3] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Springer-Verlag, Berlin, 1972, Lecture Notes in Mathematics, Vol. 304. MR0365573 (51:1825)
- [4] C. Broto and J. A. Crespo, *H-spaces with Noetherian mod two cohomology algebra*, Topology **38** (1999), no. 2, 353–386. MR1660329 (99i:55013)
- [5] W. Browder, *Torsion in H-spaces*, Ann. of Math. (2) **74** (1961), 24–51. MR0124891 (23:A2201)
- [6] N. Castellana, J. A. Crespo, and J. Scherer, *Deconstructing Hopf spaces*, Invent. Math. DOI10.1007/s00222-006-0518-8 (online).
- [7] W. Chachólski, *On the functors CW_A and PA* , Duke Math. J. **84** (1996), no. 3, 599–631. MR1408539 (97i:55023)
- [8] W. G. Dwyer, *The centralizer decomposition of BG*, Algebraic topology: new trends in localization and periodicity (Sant Feliu de Guíxols, 1994), Progr. Math., vol. 136, Birkhäuser, Basel, 1996, pp. 167–184. MR1397728 (97i:55028)
- [9] E. Dror Farjoun, *Cellular spaces, null spaces and homotopy localization*, Springer-Verlag, Berlin, 1996. MR1392221 (98f:55010)
- [10] R. Flores, *Nullification and cellularization of classifying spaces of finite groups*, to appear in Trans. Amer. Math. Soc.
- [11] R. Flores and J. Scherer, *Cellularization of classifying spaces and fusion properties of finite groups*, preprint available at <http://front.math.ucdavis.edu/math.AT/0501442>, 2005.
- [12] L. Fuchs, *Infinite abelian groups. Vol. I*, Pure and Applied Mathematics, Vol. 36, Academic Press, New York, 1970. MR0255673 (41:333)
- [13] P. Hilton, G. Mislin, and J. Roitberg, *Localization of nilpotent groups and spaces*, North-Holland Publishing Co., Amsterdam, 1975, North-Holland Mathematics Studies, No. 15, Notas de Matemática, No. 55. [Notes on Mathematics, No. 55]. MR0478146 (57:17635)
- [14] J. R. Hubbuck and R. Kane, *On π_3 of a finite H-space*, Trans. Amer. Math. Soc. **213** (1975), 99–105. MR0397724 (53:1582)
- [15] W. Meier, *Complex and real K-theory and localization*, J. Pure Appl. Algebra **14** (1979), no. 1, 59–71. MR0515487 (80b:55003)
- [16] R. J. Milgram, *The structure over the Steenrod algebra of some 2-stage Postnikov systems*, Quart. J. Math. Oxford Ser. (2) **20** (1969), 161–169. MR0248811 (40:2061)
- [17] H. Miller, *The Sullivan conjecture on maps from classifying spaces*, Ann. of Math. (2) **120** (1984), no. 1, 39–87. MR0750716 (85i:55012)
- [18] G. Mislin, *Localization with respect to K-theory*, J. Pure Appl. Algebra **10** (1977/78), no. 2, 201–213. MR0467732 (57:7585)
- [19] J.A. Neisendorfer, *Localization and connected covers of finite complexes*, The Čech centennial (Boston, MA, 1993), Contemp. Math., vol. 181, Amer. Math. Soc., Providence, RI, 1995, pp. 385–390. MR1321002 (96a:55019)
- [20] D. J. S. Robinson, *Finiteness conditions and generalized soluble groups. Part 1*, Springer-Verlag, New York, 1972, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 62. MR0332989 (48:11314)
- [21] J. L. Rodríguez and J. Scherer, *Cellular approximations using Moore spaces*, Cohomological methods in homotopy theory (Bellaterra, 1998), Progr. Math., vol. 196, Birkhäuser, Basel, 2001, pp. 357–374. MR1851263 (2003a:55024)
- [22] L. Schwartz, *Unstable modules over the Steenrod algebra and Sullivan’s fixed point set conjecture*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1994. MR1282727 (95d:55017)

- [23] R. B. Warfield, Jr., *Nilpotent groups*, Springer-Verlag, Berlin, 1976, Lecture Notes in Mathematics, Vol. 513. MR0409661 (53:13413)

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, E-08193 BELLATERRA, SPAIN

E-mail address: natalia@mat.uab.es

DEPARTAMENT DE ECONOMIA I DE HISTÒRIA ECONÒMICA, UNIVERSITAT AUTÒNOMA DE BARCELONA, E-08193 BELLATERRA, SPAIN

E-mail address: JuanAlfonso.Crespo@uab.es

Current address: Departamento de Economía, Universidad Carlos III de Madrid, E-28903 Getafe, Spain

E-mail address: jacrespo@eco.uc3m.es

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, E-08193 BELLATERRA, SPAIN

E-mail address: jscherer@mat.uab.es